



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.


We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

from the snow banks. At Horse Corral Meadow, on the way out July 21, I found the nest of a Thurber's junco. It was situated at the edge of the meadow, almost completely hidden in a clump of coarse grass. It was composed of coarse grasses, lined with finer grass,

and short hairs from mule and burro, such as might fall along the trail from clipped manes. The nest contained three downy young. The trip covered some 650 miles of wagon travel and about 200 with pack animals on mountain trails.



TYPICAL NEST OF CASSIN VIREO.

(From an enlargement of a photograph taken by R. H. Beck.)

Vireo Traits.

The above illustration typifies the nature of Cassin Vireo (*Vireo solitarius cassinii*) during its nesting season—, a confiding bird with a generous streak of curiosity running through its disposition. In June 1896 Mr. R. H. Beck found the nest represented in the illustration built in a black oak tree on the edge of a clearing, the nest being suspended only about three feet from the ground. The bird seemed not greatly disturbed by the setting up of the camera and a successful photograph resulted.

Four eggs constitutes the usual set, although I have twice found five young in a nest and one set of five eggs advanced in incubation. The nests are compactly woven and lined with light colored grasses, while the eggs are larger than those of either *Vireo huttoni* or *Vireo gilvus*. Cassin Vireo seems to possess but little fear of man and will approach very closely to the observer if he remains motionless; when the nest is disturbed the birds will alternately utter a harsh note of remonstrance and then launch into song. The location of the nest here shown was Fyffe, El Dorado Co., Cal.

C. BARLOW.